

Photometric and Spectroscopic Observations of V1280 Sco

Hiroyuki Naito¹, Sahori Mizoguchi², Akira Arai³, Masayuki Yamanaka³,
Shin-ya Narusawa¹, Kozo Sadakane² and Takashi Iijima⁴

¹*Nishi-Harima Astronomical Observatory, Sayo, Hyogo 679-5313, Japan*

²*Astronomical Institute, Osaka Kyoiku University, Kashiwara, Osaka 582-8582, Japan*

³*Hiroshima Astrophysical Science Center, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan*

⁴*Astronomical Observatory of Padova, Asiago Section, Osservatorio Astrofisico, 36012 Asiago (Vi), Italy*

ABSTRACT

Photometries of B , V , Rc , Ic , y , J , and Ks bands and low dispersion optical spectroscopic observations of Nova V1280 Sco, started soon after the outburst, are reported. We show that V1280 Sco is an Fe II nova and it is going through the historically slowest spectroscopic evolution. The rapid decline observed in the early phase was caused by formation of a dust shell. We estimate the abundances of CNO using the absorption lines on a spectrum at pre-maximum, and find over-abundances by $[C/Fe] \sim 1.4$, $[N/Fe] > 2.0$ and $[O/Fe] \sim 1.1$.

Key words: photometry, spectroscopy, nova: individual (V1280 Sco), nova:dust nova

1 INTRODUCTION

V1280 Sco (Nova Scorpii 2007 No. 1) is a classical nova which was discovered by Y. Nakamura and Y. Sakurai on the same night (Feb. 4, 2007) at 9-th visual magnitude (Yamaoka et al. 2007a). Its early spectra were taken by some observers and it was classified as an Fe II type nova (e.g. Naito & Narusawa 2007, Munari et al. 2007, Kuncarayakti et al. 2008), where the Fe II type nova spectrum is described in Williams (1992). Yamaoka et al. (2007b) reported that the spectrum at the pre-maximum stage looked like an early-type supergiant dominated by absorption lines of hydrogen, iron and other metals. It showed a somewhat slow rise to maximum of about 3.7 in the V band on Feb. 16 (12 days after the discovery), and became a naked eye nova since V382 Vul and V1494 Aql recorded in 1999. It is also remarkable in that V1280 Sco formed a dust shell in the early phase. After maximum, it faded slowly for about 12 days, then it declined rapidly in the visual region by the obscuration due to dust shell(s), the formation of which was directly detected by Chesneau et al. (2008) from near-IR and mid-IR observations by the VLTI at ESO. Das et al. (2008) presented near-infrared studies of V1280 Sco and suggested that the dust is in clumpy shells.

2 PHOTOMETRY

We performed B , V , Rc , Ic and y photometries using a 51-cm reflector at Osaka Kyoiku University and J and Ks photometries using the 1.5-m KANATA telescope at Higashi-Hiroshima Observatory. The light curves in V , J and Ks bands are shown in Fig. 1. Concerning the visual light,




Figure 1. Light curves in V (filled circle), J (open triangle) and Ks (open square) bands.

within two weeks after maximum, V1280 Sco started a rapid decline until mid-May when a re-brightening started. On the other hand, the brightness in the Ks band stayed at nearly the same brightness ($Ks \sim 4$) through these days. These results imply that V1280 Sco formed thick dust shell(s), which are consistent with the results of Chesneau et al. (2008) and Das et al. (2008). When the first re-brightening in the V band occurred in the latter half of May 2007, near IR magnitudes had brightened simultaneously. We guess that another H-burning event had happened on the WD at that time. The light curves in visual show the second re-brightening from




Figure 2. Spectral series of V1280 Sco from Feb. 5, 2007 to Jun. 10, 2007. These spectra are corrected for reddening based on the preliminary $E(B-V)$ value of 0.38 determined from photometries near maximum.




Figure 3. The last spectrum taken on Sep. 9, 2008. The reddening is not corrected.

the beginning of Aug., which is interpreted that the obscuring dust became thinner gradually.

3 SPECTROSCOPY

We performed optical spectroscopic observations at Nishi-Harima Astronomical Observatory (NHAO), using the 2.0-m NAYUTA telescope equipped with the MALLS (Medium And Low dispersion Long slit Spectrograph). We obtained low dispersion spectra covering the range 420–680 nm with a resolution of $\lambda / \Delta \lambda \cong 1200$. Fig. 2 shows selected spectra of V1280 Sco observed at NHAO.

The first spectrum taken on Feb. 5 displays a smooth continuum with the Balmer and Fe II lines in emission, accompanied by P Cygni absorptions. The expansion velocity estimated from the blue-shift of the absorption component of $H\alpha$ was 480 km s^{-1} . This spectrum shows that V1280 Sco

can be classified as an Fe II type nova. On Feb. 14, two days before maximum, the spectrum showed a dramatic change. It apparently became an early A-type supergiant. From an analysis of absorption lines observed on this spectrum, we estimate the abundances of CNO contained of this object, and found over-abundances by $[C/Fe] \sim 1.4$, $[N/Fe] > 2.0$ and $[O/Fe] \sim 1.1$. The analysis is based on an assumption that the atmospheric parameters of V1280 Sco on Feb. 14 could be approximated by those of Deneb (A2Ia).

The spectral evolution after maximum luminosity was very slow and strong forbidden lines of [O I] $\lambda 5577$, [O I] $\lambda 6300, 6363$ were observed on Mar. 7. The ratio of [O I] $\lambda 5577$ to [O I] $\lambda 6300$ is so large that we can infer the electron density in the ejecta was still high because the ejected mass is large and, at the same time, the expansion velocity is low. When the optical decline started by the dust formation from March to mid-May, the continuum flux was too weak to be detected and at the same time the $H\beta$ emission also was not detected. However, during the re-brightening phase, $H\beta$ line reappeared. On the same spectrum a P Cygni absorption line having a velocity of $2,000 \text{ km s}^{-1}$ at $H\alpha$ was shown. These may have been connected with a new mass ejection on the re-brightening. It is caused by the second large outflow triggered by a new H-burning event on the WD. Until June 10, no emission line of He I, He II, or other highly ionized ions such as C III or N III was observed.

On Feb. 18, 2008, the He I $\lambda 6678$ appeared and the blue continuum was strong. This indicated that the temperature was higher than in 2007, but the temperature was not so high as to excite the He II lines. A higher S/N spectrum observed on Jul. 8 showed P-Cyg profiles in the Balmer lines, indicating the gas outflow caused by the continueing H burning. Even on our last spectrum taken on Sep. 9, 2008, the forbidden lines of [O III] $\lambda 4959, 5007$ were not seen, thus V1280 Sco has not entered the nebular phase yet (Fig. 3).

4 SUMMARY

From our observations, it is clear that V1280 Sco is a very unique nova in history. GQ Mus had the H burning turn-off time of about 3,000 days (Hachisu et al. 2008), and V723 Cas took 540 days before entering the nebular phase (Iijima 2006). V1280 Sco takes longer time than V723 Cas to enter the nebular phase, and it may rewrite the record of GQ Mus. Follow-up observations are needed to probe the nature of this very interesting nova.

REFERENCES

- Chesneau, O. et al., 2008, A&A, 487, 223
- Das, R. K. et al., 2008, astro-ph/0809.4338
- Hachisu, I., Kato, M., Cassatella, A., 2008, astro-ph/0806.4253
- Iijima, T., 2006, A&A, 451, 563
- Kuncarayakti, H., Kristyowati, D., Kunjaya, C., 2008, Ap&SS, 314, 209
- Munari, U. et al., 2007, CBET 852
- Naito H., Narusawa S., 2007, IAC Circ. 8803
- Williams, R. E., 1992, AJ, 104, 725
- Yamaoka, H., et al., 2007a, IAU Circ. 8803
- Yamaoka, H., Fujii, M., Naito, H., 2007b, IAU Circ. 8807